

Peroxidase Blocking Solution

<u>Cat No.</u>	<u>Quantity</u>
10-0056	100 mL Ready-To-Use

Intended Use	For In Vitro Diagnostic Use. This product is intended for use to block endogenous peroxidase on formalin-fixed, paraffin embedded (FFPE) tissues during the pretreatment process of the immunohistochemical (IHC) staining procedure.
Reagents Supplied	One 100 mL bottle of Ready-To-Use Peroxidase Blocking Solution containing 3% hydrogen peroxide in reagent water.
Summary And Explanation	This product strongly inhibits endogenous peroxidase in formalin-fixed, paraffin-embedded tissue sections.
Procedure	The blocking can be done 1) after rehydration and before antigen retrieval, 2) after antigen retrieval and before primary antibody incubation, 3) after primary antibody incubation, or 4) after secondary antibody incubation. Incubate the slides for 10 -15 minutes at room temperature. Rinse with PBS 3 times.
Storage	Store at 2-8°C and protect from sunlight. Do not shake bottle. All performance claims are void after the expiration date.
Materials Required But Not Supplied	FFPE tissue section PBS
Precautions	For professional users only. Safety data sheet (MSDS) available for professional users upon request.

Symbols

				
Catalog No.	Batch No.	In Vitro Diagnostic Use	Temperature Range	Use By

