

Mouse Monoclonal anti-Cytokeratin 19, Clone RCK108

60-0112; 60-0112-7	6 mL; 7 mL predilute Antibody, Ready-To-Use
61-0112-2	0.2 mL Concentrate Antibody
Isotype	IgG1
Concentration	See container label

Intended Use

For In Vitro Diagnostic Use.

This product is used to qualitatively detect Cytokeratin 19 in normal and neoplastic formalin fixed, paraffin embedded (FFPE) tissue sections in immunohistochemical (IHC) detection methodology. Interpretation must be made within the context of the patient's clinical history and other diagnostic test by a qualified pathologist.

Description

The antibody labels many types of normal and neoplastic simple and non-keratinizing epithelia, including ductal and glandular epithelia. It is a useful tool for the identification of epithelial tumors.

Reagent provided

This antibody is diluted in 10 mM phosphate buffered saline (PBS), pH 7.2 containing 1% bovine serum albumin (BSA) and 0.09% sodium azide (NaN₃) as antimicrobial agent.

Precautions

For professional users.

Proper handling of this product as with any product derived from biological sources according to local and applicable regulations.

Sodium azide (NaN₃) is a toxic chemical. The concentration in this product is not classified as hazardous, however, the build-ups of NaN₃ may react with lead and copper plumbing to form highly explosive metal azides. Flush the disposed reagent with large volume of water to prevent azide build-up.

Usage

Dilution

60-0112; 60-0112-7: Ready-To-Use

61-0112-2: Dilute 1:50-100 before use when using Acu-Stain™ detection system. Optimum dilution factor may vary depending on the specimen and preparation process and should be determined by each individual investigator.

Staining procedure

Incubate this antibody with tissue section for 30-60 minutes at room temperature. Follow the instructions from the selected detection system.

Positive control tissue

Tonsil, GI track

Epitope retrieval

Proteinase K

Staining pattern

Cytoplasm

Storage

Store at 2-8°C.

References

1. Kwaspen F, et al. Histopathology. 1997 Dec;31(6):503-16.
2. Moll R. Subcell Biochem. 1998;31:205-62.

Symbols

				
Catalog No.	Batch No.	In Vitro Diagnostic Use	Temperature Range	Use By