

Mouse Monoclonal anti-MUM1, Clone MUM1p

60-0136; 60-0136-7 6 mL; 7 mL predilute Antibody, Ready-To-Use
 61-0136; 61-0136-2; 61-0136-5 1 mL; 0.2 mL; 0.5 mL Concentrate Antibody
 Isotype IgG1
 Concentration See container label

Intended Use For In Vitro Diagnostic Use.
 This product is used to qualitatively detect MUM1 in normal and neoplastic formalin fixed paraffin embedded (FFPE) tissue sections in immunohistochemical (IHC) detection methodology. Interpretation must be made within the context of the patient's clinical history and other diagnostic test by a qualified pathologist.

Description The antibody labels the MUM1 protein, which is expressed in a subset of B cells in the light zone of the germinal center, plasma cells, activated T cell, a wide spectrum of related hematolymphoid neoplasms. The antibody is useful for the subclassification of lymphoid malignancies.

Reagent provided This antibody is diluted in 10 mM phosphate buffered saline (PBS), pH 7.2 containing 1% bovine serum albumin (BSA) and 0.09% sodium azide (NaN₃) as antimicrobial agent.

Precautions For professional users.
 Proper handling of this product as with any product derived from biological sources according to local and applicable regulations.
 Sodium azide is a toxic chemical. The concentration in this product is not classified as hazardous, however, the build-up of NaN₃ may react with lead and copper plumbing to form highly explosive metal azides. Flush the disposed reagent with large volume of water to prevent azide build-up.

Usage
Dilution 60-0136; 60-0136-7: Ready-To-Use
 61-0136; 61-0136-2; 61-0136-5: Dilute 1:50 to 1:100 before use when using Acu-Stain™ detection system. Optimum dilution factor may vary depending on the specimen and preparation process and should be determined by each individual investigator.

Staining procedure Incubate this antibody with tissue section for 30-60 minutes at room temperature. Follow the instructions from the selected detection system.

Positive control tissue Tonsil or Myeloma

Epitope retrieval HIER, Citrate pH 6 or Tris pH 9

Staining pattern Nucleus

Storage Store at 2-8°C.

- References**
1. Falini B, et al. Blood. 2000 Mar 15;95(6):2084-92.
 2. Natkunam Y, et al. Mod Pathol. 2001 Jul;14(7):686-94.

Symbols

 Catalog No.	 Batch No.	 In Vitro Diagnostic Use	 Temperature Range	 Use By
--	--	---	--	---